



An Introduction to Polymerase Chain Reaction (PCR)

Ameer Effat M. Elfarash

Dept. of Genetics
Fac. of Agriculture, Assiut Univ.
aelfarash@aun.edu.eg

Introduction

- **The technique was invented by Dr. Kary Mullis, 1986**
- **for which he received the Nobel Prize in Chemistry in 1993.**



PCR Achieves Fame and Fortune --becomes standard in molecular biology tool box--





The Molecule of the Year

RUTH LEVY GUYER AND DANIEL E. KOSHLAND, JR.


Science HAS SELECTED THE POLYMERASE CHAIN REACTION AS the major scientific development of 1989 and has chosen for its first "Molecule of the Year" the DNA polymerase molecule that drives the reaction. The list from which the polymerase chain reaction (PCR) was chosen included an impressive array of accomplishments in many areas of science and technology; additional kudos are therefore conferred below to 17 of the other big "stories" that made 1989 an exciting year for scientists and for followers and beneficiaries of science. Although the PCR procedure was introduced several years ago, use of the technique truly burgeoned in 1989; in much the same way, the full potentials of many of the interesting "runner-up" scientific achievements of this year are likely to be realized sometime in the years to come.



DNA Structure




DNA Replication



Polymerase Chain Reaction (PCR)


- PCR is a technique which is used to amplify the number of copies of a **specific region of DNA**, (usually fewer than 3000 base pairs) in order to produce enough DNA to be adequately tested.
- **Millions** of copies of a segment of DNA can be made within a few hours
- As a result, it now becomes possible to analyze and characterize the DNA.

● DNA amplification by PCR (overview)




PCR Cycle


- **Each cycle (Round) of PCR contains 3 steps:**
 - 1- Denaturation**
 - 2- Primer annealing**
 - 3- Primer extension**
- **The cycle usually repeated for 25 – 40 times.**



Programming the Thermocycler





Reaction buffer
DNA template
PCR primers
Taq DNA pol




Denature Template

Anneal Primers


DNA Synthesis





PCR



PCR





PCR










PCR








PCR





PCR




PCR





PCR



Time



Fragments of defined length





Movie

What do we need for PCR?



PCR tube



THERMOCYCLER

متطلبات تقنيه تفاعل البوليمر المتسلسل (PCR)

DNA Sample



Primers



CCGAATGGGATGC

GGCTTACCCTACG

نوعان :

• أمامي (Forward).

• خلفي (Reverse).

وهي تتابع من القواعد النيتروجينية في شريط واحد
قصير (20-25 b) مكمل لبداية الجزء المراد تضخيمه في الـ
DNA.

Taq polymerase



- مستخرج من سلالة بكتيرية تسمى *Thermus aquaticus* التي تعيش في المياه الحارة.

- لا يتأثر بدرجات الحرارة المرتفعة.
- درجة الحرارة المثلى له ٧٢ °م.

First reports using DNA polymerase from *Thermus aquaticus* was at (1988)

dNTPs



Adenine

أدينين



Thymine

ثايمين



Guanine

جوانين



Cytosine

سايتوسين






PCR Buffer 10x



Distilled Water

PCR Procedure

All the required components are inserted into an Eppendorf tube




Template DNA

salts (ions)

dNTPs

DNA Polymerase

Primers



PROCEDURE

PCR




Agarose gel electrophoresis




3-4 hours

Reliable PCR from Every Sample



The final product



UV visualisation

PROCEDURE

- DNA (Template).
- Forward primer
- Reverse primer
- dNTP's
- *Taq* DNA Polymerase
- Buffer
- H₂O

